





Date	Focus	Platform	Post	Link
11-May	Hospitals	Facebook	<p>About 900,000 people in the U.S. experience blood clots each year, and about half of these clots occur within three months of a hospital stay or surgery. Blood clots are a leading cause of preventable hospital death in the United States. If you are going to be in the hospital or have surgery, make sure all of your doctors know your risk for blood clots and ask for a prevention plan before, during, and after your stay. Learn more during National Hospital Week: http://bit.ly/2qADWRT #stoptheclot #spreadtheword #sharetostoptheclot #hospitalweek</p>	https://www.stoptheclot.org/spreadtheword/hospitalization/
11-May	Hospitals	Twitter	<p>A hospital stay and surgery are risk factors for blood clots. Work with your doctors to have a prevention plan before, during and after your stay. Learn more during National Hospital Week: http://bit.ly/2qADWRT #stoptheclot #spreadtheword #hospitalweek</p>	https://www.stoptheclot.org/spreadtheword/hospitalization/

11-May	Hospitals	Instagram	About 900,000 people in the U.S. experience blood clots each year, and about half of these clots occur within three months of a hospital stay or surgery. Blood clots are a leading cause of preventable hospital death in the United States. If you are going to be in the hospital or have surgery, make sure all of your doctors know your risk for blood clots and ask for a prevention plan before, during, and after your stay. Visit the link in our bio profile @stoptheclot to learn more or type www.stoptheclot.org/spreadtheword into your Internet browser.	N/A
12-May	Hospitals	Facebook	If you are confined to a bed or wheelchair for long periods of time due to a hospital stay, injury, or serious illness, you are at increased risk for a blood clot. Click here to download a printable Infographic that can help you, along with your doctor, determine your blood clot risk prior to a hospital stay, or surgery, and work together to develop a blood clot prevention plan: http://bit.ly/2ECL875 . Also, be sure to know the signs and symptoms of blood clots, and let your doctor know right away if you experience any of them. #stoptheclot #spreadtheword #sharetostoptheclot #hospitalweek	https://www.stoptheclot.org/spreadtheword/wp-content/uploads/2018/01/Hospitalization-InfographicPRINT.pdf
12-May	Hospitals	Twitter	If you are confined to a bed or wheelchair for long periods of time due to a hospital stay, injury, or serious illness, you are at increased risk for a blood clot. Have a prevention plan and know the signs: http://bit.ly/2ECL875 #hospitalweek	https://www.stoptheclot.org/spreadtheword/wp-content/uploads/2018/01/Hospitalization-InfographicPRINT.pdf

Image	Reach	Reactions	Clicks	Impressions
 <p>STOP THE CLOT SPREAD THE WORD™</p> <p>Heading to the Hospital? Get Better. Don't Get a Blood Clot.</p> <p>For more information, visit stoptheclot.org/spreadtheword</p> <p>National Blood Clot Alliance CDC CENTERS FOR DISEASE CONTROL AND PREVENTION</p>	2,863	51	25	N/A
 <p>STOP THE CLOT SPREAD THE WORD™</p> <p>Heading to the Hospital? Get Better. Don't Get a Blood Clot.</p> <p>For more information, visit stoptheclot.org/spreadtheword</p> <p>National Blood Clot Alliance CDC CENTERS FOR DISEASE CONTROL AND PREVENTION</p>	N/A	N/A	7	977

(Same as Facebook)	696	N/A	2	696
	2,784	45	25	N/A
	N/A	N/A	7	898

Total Engagement

N/A

15

N/A

N/A

11

N/A
N/A
28

